



Therya

ISSN: 2007-3364

Asociación Mexicana de Mastozoología A. C.

Sánchez-Montes, Sokani; Cabrera-Garrido, Martín Yair; Ríos-Muñoz, César A.; Lira-Olguín, Ali Zeltzin; Acosta-Gutiérrez, Roxana; Mata-Galindo, Mario; Hernández-Vilchis, Kevin; Navarrete-Sotelo, D. Melissa; Colunga-Salas, Pablo; León-Paniagua, Livia; Becker, Ingeborg
Detection of *Bartonella* and *Rickettsia* in small mammals and their ectoparasites in México

Therya, vol. 10, no. 2, 2019, pp. 69-79
Asociación Mexicana de Mastozoología A. C.

DOI: 10.12933/therya-19-722

Available in: <http://www.redalyc.org/articulo.oa?id=402362668002>

- ▶ How to cite
- ▶ Complete issue
- ▶ More information about this article
- ▶ Journal's webpage in redalyc.org

The logo for Redalyc, featuring the text 'redalyc.org' with a stylized red and black graphic element.

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Detection of *Bartonella* and *Rickettsia* in small mammals and their ectoparasites in México

SOKANI SÁNCHEZ-MONTES¹, MARTÍN YAIR CABRERA-GARRIDO², CÉSAR A. RÍOS-MUÑOZ^{1,3}, ALI ZELTZIN LIRA-OLGUIN²; ROXANA ACOSTA-GUTIÉRREZ², MARIO MATA-GALINDO¹, KEVIN HERNÁNDEZ-VILCHIS¹, D. MELISSA NAVARRETE-SOTELO¹, PABLO COLUNGA-SALAS^{1,2}, LIVIA LEÓN-PANIAGUA² AND INGEBORG BECKER^{1*}

¹ Centro de Medicina Tropical, Unidad de Medicina Experimental, Facultad de Medicina. Universidad Nacional Autónoma de México. Dr. Balmis 148, CP. 06726, Ciudad de México. México. Email: sok10108@gmail.com (SSM), rmunoz98@gmail.com (CARM), mata316@ciencias.unam.mx (MMG), chopa95@ciencias.unam.mx (KHV), dmel1231@gmail.com (MNS), colungasalas@gmail.com (PCS), becker@servidor.unam.mx (IB).

² Museo de Zoología "Alfonso L. Herrera", Departamento de Biología Evolutiva, Facultad de Ciencias, Universidad Nacional Autónoma de México. Avenida Universidad 3000, CP. 04510, Ciudad de México México. Email: bmw_mark@comunidad.unam.mx (MYCG), alizeltzin@ciencias.unam.mx (AZLO), roxana_a2003@yahoo.com.mx (RAG), colungasalas@gmail.com (PCS), llp@ciencias.unam.mx (LLP).

³ Laboratorio de Arqueozoología, Subdirección de Laboratorios y Apoyo Académico, Instituto Nacional de Antropología e Historia. Moneda 16, CP. 06060, Ciudad de México. México. Email: rmunoz98@gmail.com (CARM).

* Corresponding author.

Fleas and sucking lice are important vectors of multiple pathogens causing major epidemics worldwide. However these insects are vectors of a wide range of largely understudied and unattended pathogens, especially several species of bacteria's of the genera *Bartonella* and *Rickettsia*. For this reason the aim of the present work was to identify the presence and diversity of *Bartonella* and *Rickettsia* species in endemic murine typhus foci in Hidalgo, México. A cross-sectional study was carried out to collect small mammals and their associated ectoparasites during October, 2014. Samples of liver and ear of hosts, and ectoparasites were fixed in absolute ethanol and examined to identify the presence of *Bartonella* and *Rickettsia* DNA by the amplification of specific fragments of the *gltA* and *ompB* genes using conventional PCR. The recovered sequences were compared with those deposited in GenBank, and phylogenetic analyzes were carried out to identify the position of the pathogens detected with respect to the valid species previously reported worldwide. A total of 47 fleas and 172 sucking lice, belonging to five families (Ceratothyllidae, Leptopsyllidae, Ctenophtalmidae, Hoplopleuridae, Polyplacidae) and related to six species were collected from 40 rodents of four species and one shrew. Only four hosts (two *P. beatae*, and two *R. norvegicus*) were positive to *Bartonella elizabethae*, *Bartonella vinsonii* and *Rickettsia typhi*. In the case of ectoparasites, 23 specimens of two flea species (*Peromyscopsylla hesperomys* and *Plusaetis mathesoni*) tested positive for *B. vinsonii*. No evidence of *Bartonella* or *Rickettsia* was detected in any lice. Our findings represent the first record of *Bartonella elizabethae* a confirmed zoonotic pathogen causing endocarditis in México and several new associations of *Bartonella* with Mexican flea species, which highlight the importance of the establishment of active entomological surveillance in wildlife.

Las pulgas y los piojos son vectores de patógenos causantes de epidemias de importancia histórica. Sin embargo, estos insectos son vectores de una amplia gama de patógenos poco estudiados y no atendidos, especialmente varias especies de bacterias de los géneros *Bartonella* y *Rickettsia*. Por este motivo, el objetivo del presente trabajo fue identificar la presencia y diversidad de las especies de *Bartonella* y *Rickettsia* en un foco de tifus murino en el estado de Hidalgo, México. Se realizó un estudio transversal para recolectar hospederos y sus ectoparásitos durante octubre de 2014. Las muestras de hígado y oreja de los hospederos y los ectoparásitos se fijaron en etanol absoluto y se examinaron para identificar la presencia de ADN de *Bartonella* y *Rickettsia* mediante la extracción de DNA y amplificación de fragmentos específicos de los genes *gltA* y *ompB*. Las secuencias obtenidas fueron comparadas con aquellas depositadas en GenBank y se realizaron análisis filogenéticos para identificar la posición de los patógenos detectados respecto a las especies válidas previamente reportadas a nivel mundial. Se recolectaron un total de 47 pulgas y 172 piojos chupadores, pertenecientes a seis especies de cinco familias (Ceratothyllidae, Leptopsyllidae, Ctenophtalmidae, Hoplopleuridae, Polyplacidae) asociados con 40 roedores de cuatro especies y una musaraña. Sólo cuatro hospederos (dos *P. beatae*, y dos *R. norvegicus*) resultaron positivos para *Bartonella elizabethae*, *Bartonella vinsonii* y *Rickettsia typhi*. En el caso de los ectoparásitos, 23 ejemplares de dos especies de pulgas (*Peromyscopsylla hesperomys* y *Plusaetis mathesoni*) fueron positivos para *B. vinsonii*. No se detectó evidencia de ninguno de los dos patógenos en los piojos analizados. Nuestros hallazgos representan el primer registro de *Bartonella elizabethae*, un patógeno zoonótico confirmado que causa endocarditis en México y varias asociaciones nuevas de *Bartonella* con especies de pulgas mexicanas, lo cual resalta la necesidad de implementar vigilancia entomológica activa para el monitoreo de estos patógenos en animales silvestres.

Keywords: *Bartonella elizabethae*; emerging diseases; *Rickettsia typhi*; small mammals; vectors.

© 2019 Asociación Mexicana de Mastozoología, www.mastozoologiamexicana.org

Introduction

Fleas and sucking lice are important vectors of multiple pathogens causing major epidemics worldwide, such as plague (*Yersinia pestis*) and epidemic typhus (*Rickettsia prowazekii*). Despite the historical importance of both diseases, this group of ectoparasites has been little studied

with respect to other vectors such as mosquitoes or ticks (Gillespie *et al.* 2009; Bitam *et al.* 2010; Eisen and Gage 2012). However, these groups of insects are hosts for a wide range of largely understudied pathogens, especially several species of bacteria of the genera *Bartonella* and *Rickettsia* (Bitam *et al.* 2010). The genus *Bartonella* includes

at least 33 species of Gram-negative, intracellular and slow-growing coccobacilli with complex life cycles including multiple vertebrate hosts and vectors, such as *B. elizabethae* and *B. vinsonii arupensis*, declared pathogens causing endocarditis in humans and dogs (Breitschwerdt and Kordick 2000; Tsai et al. 2011; Kosoy et al. 2012; Regier et al. 2016). On the other hand, *Rickettsia* encompasses 26 species of obligate intracellular bacteria which are transmitted by different groups of hematophagous arthropods such as ticks, lice and fleas (Fournier and Raoult 2009; Merhej et al. 2014). *Rickettsia* species are classified into four groups, two of which are pathogens for man: members of the Spotted Fever group [SGF] (*R. conorii*, *R. massiliae*, *R. rickettsii* and *R. parkeri*) and Typhus group [TG] (*R. prowazekii* and *R. typhi*), this latter group is transmitted exclusively by lice and fleas,

which cause epidemic and murine typhus (Fournier et al. 2003; Fournier and Raoult 2009).

In recent decades with the advent of molecular biology techniques, the number of species or strains of both bacteria genera has increased exponentially (Merhej et al. 2014; Regier et al. 2016). Particularly, fleas and sucking lice associated with rodents are the groups in which more studies have focused for the detection of pathogens, with the identification of 16 validated species of *Bartonella*, nine of *Rickettsia* and more than 17 new lineages near to several validated taxa (but which require isolation for formal identification) for both genera, associated with 45 flea species and seven sucking lice which are also associated with 42 species of rodents in 24 countries around the world (Table 1).

Table 1. *Bartonella* and *Rickettsia* species detected in fleas and sucking lice associated with rodents worldwide

Bacteria species	Flea	Host	Country	References	
<i>B. birtlesii</i>	<i>Ctenophthalmus andorrensis catalanensis</i>	<i>Apodemus sylvaticus</i>	Spain	Cevidane et al. 2017	
	<i>Leptopsylla taschenbergi amitina</i>	<i>A. sylvaticus</i>	Spain	Cevidane et al. 2017	
<i>B. coopersplainsensis</i>	<i>Stephanocircus pectinipes</i>	<i>Rattus fuscipes</i>	Australia	Kaewmongkol et al. 2011	
<i>B. doshiae</i>	<i>Xenopsylla cheopis</i>	<i>Rattus</i> sp.	Afghanistan	Marie et al. 2006	
<i>B. elizabethae</i>	<i>Leptopsylla segnis</i>	<i>Mus spretus</i>	Algeria	Bitam et al. 2012	
	<i>Synosternus cleopatrae</i>	<i>Gerbillus pyramidum</i>	Israel	Morick et al. 2010	
	<i>Synopsyllus fonquerniei</i>	<i>Rattus rattus</i>	Madagascar	Brook et al. 2017	
	<i>X. cheopis</i>	<i>Rattus norvegicus</i>	Algeria	Bitam et al. 2012	
			USA	Frye et al. 2015	
			<i>R. rattus</i>	Algeria	Bitam et al. 2012
			<i>Rattus tanezumi</i>	Indonesia	Winoto et al. 2005
		<i>Rattus</i> sp.	Afghanistan	Marie et al. 2006	
			Nigeria	Kamani et al. 2013	
<i>B. grahamii</i>	<i>Ctenophthalmus agyrtes</i>	ND	Lithuania	Lipatova et al. 2015	
	<i>Ct. andorrensis catalanensis</i>	<i>A. sylvaticus</i>	Spain	Cevidane et al. 2017	
	<i>Ctenophthalmus nobilis</i>	<i>Myodes glareolus</i>	England	Bown et al. 2004	
	<i>Megabothris turbidus</i>	ND	Lithuania	Lipatova et al. 2015	
	<i>Megabothris walkeri</i>	ND	Lithuania	Lipatova et al. 2015	
	<i>Sy. cleopatrae</i>	ND	Israel	Rzotkiewicz et al. 2015	
	<i>Xenopsylla ramesis</i>	ND	Israel	Rzotkiewicz et al. 2015	
<i>B. henselae</i>	<i>X. ramesis</i>	ND	Israel	Rzotkiewicz et al. 2015	
		<i>Meriones tristrami</i>	Israel	Morick et al. 2010	
<i>B. koehlerae</i>	<i>Xenopsylla gerbilli</i>	<i>Meriones lybicus</i>	Afghanistan	Marie et al. 2006	
<i>B. phoceensis</i>	<i>X. cheopis</i>	<i>R. tanezumi</i>	Indonesia	Winoto et al. 2005	
<i>B. queenslandensis</i>	<i>X. cheopis</i>	<i>Rattus</i> sp.	Thailand	Klangthong et al. 2015	
<i>B. quintana</i>	<i>X. gerbilli</i>	<i>Meriones lybicus</i>	Afghanistan	Marie et al. 2006	
<i>B. rattaustaliani</i>	<i>Stephanocircus dasyure</i>	<i>R. fuscipes</i>	Australia	Kaewmongkol et al. 2011	
<i>B. rattimassiliensis</i>	<i>X. cheopis</i>	<i>R. tanezumi</i>	Indonesia	Winoto et al. 2005	
<i>B. rochalimae</i>	<i>X. cheopis</i>	<i>R. norvegicus</i>	USA	Frye et al. 2015	
<i>B. taylorii</i>	<i>Ct. agyrtes</i>	ND	Lithuania	Lipatova et al. 2015	
	<i>Ct. andorrensis catalanensis</i>	<i>A. sylvaticus</i> , <i>C. russula</i> , <i>M. spretus</i>	Spain	Cevidane et al. 2017	
	<i>Ct. nobilis</i>	<i>M. glareolus</i>	England	Bown et al. 2004	
	<i>Ctenophthalmus uncinatus</i>	ND	Lithuania	Lipatova et al. 2015	
	<i>Hystrichopsylla talpae</i>	ND	Lithuania	Lipatova et al. 2015	

Continue...

Bacteria species	Sucking lice	Host	Country	References
	<i>L. taschenbergi amitina</i>	<i>A. sylvaticus</i>	Spain	Cevidaneš et al. 2017
	<i>M. turbidus</i>	ND	Lithuania	Lipatova et al. 2015
	<i>M. walkeri</i>	ND	Lithuania	Lipatova et al. 2015
	<i>X. gerbilli</i>	<i>M. lybicus</i>	Afghanistan	Marie et al. 2006
<i>B. tribocorum</i>	<i>Ctenophthalmus</i> sp.	ND	Nigeria	Kamani et al. 2013
	<i>X. cheopis</i>	<i>R. norvergicus</i>	USA	Reeves et al. 2007a; Frye et al. 2015
		<i>R. rattus</i>	Algeria	Bitam et al. 2012
		<i>R. tanezumi flavipectus</i>	China	Li et al. 2007
		<i>Rattus</i> sp.	Thailand	Klangthong et al. 2015
<i>B. vinsonii</i>	<i>Polygenis bohlsi bohlsi</i>	<i>Thrichomys fosteri</i>	Brazil	de Sousa et al. 2018
	<i>Polygenis gwyni</i>	<i>Sigmodon hispidus</i>	USA	Abbot et al. 2007
<i>B. vinsonii arupensis</i>	<i>Malareus sinomus</i>	<i>Peromyscus eremicus</i>	México	Zapata-Valdés et al. 2018
	<i>Orchopeas leucopus</i>	<i>P. eremicus</i>		Fernández-González et al. 2016
		<i>Peromyscus leucopus, Peromyscus maniculatus</i>		Fernández-González et al. 2016
	<i>Pleochaetis exilis</i>	<i>Onychomys torridus</i>		Zapata-Valdés et al. 2018
<i>B. vinsonii vinsonii</i>	<i>Ctenophthalmus pseudagyrtis</i>	<i>Microtus</i> sp.	USA	Reeves et al. 2007a
	<i>Meringis parkeri</i>	<i>Onychomys arenicola, Onychomys leucogaster</i>	México	Fernández-González et al. 2016
	<i>Orchopeas sexdentatus</i>	<i>Neotoma albigula</i>	México	Fernández-González et al. 2016
	<i>Pleochaetis exilis</i>	<i>N. albigula, O. arenicola, O. leucogaster, P. maniculatus</i>	México	Fernández-González et al. 2016
<i>B. washoensis</i>	<i>Orchopeas hirsuta</i>	<i>Cynomys</i> sp.	USA	Stevenson et al. 2003; Reeves et al. 2007b
		<i>Cynomys ludovicianus</i>	México	Zapata-Valdés et al. 2018
	<i>Orchopeas howardi</i>	<i>Sciurus carolinensis</i>	USA	Durden et al. 2004
	<i>Oropsylla montana</i>	<i>Otospermophilus beecheyi</i>	USA	Osikowicz et al. 2016
	<i>Pulex</i> sp.	<i>C. ludovicianus</i>	México	Fernández-González et al. 2016
	<i>Thrassis fatus</i>	<i>Cynomys</i> sp.	USA	Reeves et al. 2007b
<i>Bartonella</i> near <i>birtlesii</i>	<i>O. howardi</i>	<i>S. carolinensis</i>	USA	Reeves et al. 2005b
<i>Bartonella</i> near <i>clarridgeiae</i>	<i>Ctenophthalmus lushuiensis</i>	<i>Eothenomys</i> sp.	China	Li et al. 2007
	<i>L. segnis</i>	<i>R. rattus</i>	Egypt	Loftis et al. 2006
	<i>P. gwyni</i>	<i>S. hispidus</i>	USA	Abbot et al. 2007
<i>Bartonella</i> near <i>doshiae</i>	<i>Ct. andorrensis catalanensis</i>	<i>A. sylvaticus</i>	Spain	Cevidaneš et al. 2017
	<i>L. taschenbergi amitina</i>	<i>A. sylvaticus</i>	Spain	Cevidaneš et al. 2017
<i>Bartonella</i> near <i>elizabethae</i>	<i>Ct. andorrensis catalanensis</i>	<i>A. sylvaticus</i>	Spain	Cevidaneš et al. 2017
	<i>Leptopsylla algira</i>	ND	Israel	Rzotkiewicz et al. 2015
		<i>Mus musculus</i>	Israel	Morick et al. 2010
	<i>L. taschenbergi amitina</i>	<i>A. sylvaticus</i>	Spain	Cevidaneš et al. 2017
	<i>Ornithophaga</i> sp.	<i>M. spretus</i>	Portugal	De Sousa et al. 2006
	<i>Stenoponia tripectinata</i>	<i>M. spretus</i>	Portugal	De Sousa et al. 2006
		<i>R. rattus</i>	Portugal	De Sousa et al. 2006
	<i>Sy. cleopatrae</i>	ND	Israel	Rzotkiewicz et al. 2015
		<i>G. pyramidum</i>	Israel	Morick et al. 2010
	<i>X. cheopis</i>	<i>Rattus</i> sp.	Thailand	Klangthong et al. 2015
	<i>X. ramesis</i>	ND	Israel	Rzotkiewicz et al. 2015
<i>Bartonella</i> near <i>grahamii</i>	<i>Meringis altipecten</i>	<i>O. arenicola, O. leucogaster, Dipodomys merriami</i>	México	Fernández-González et al. 2016
	<i>Meringis arachis</i>	<i>O. arenicola, O. leucogaster, D. merriami</i>	México	Fernández-González et al. 2016
	<i>M. parkeri</i>	<i>O. arenicola, O. leucogaster, D. merriami</i>	México	Fernández-González et al. 2016
	<i>Nosopsyllus fasciatus</i>	<i>Rattus surifer</i>	Thai-Myanmar Border	Parola et al. 2003

Continue...

Bacteria species	Sucking lice	Host	Country	References
	<i>P. exilis</i>	<i>O. arenicola</i> , <i>O. leucogaster</i>	México	Fernández-González <i>et al.</i> 2016
	<i>Sy. cleopatrae</i>	<i>Meriones sacramenti</i>	Israel	Morick <i>et al.</i> 2010
	<i>X. ramesis</i>	ND	Israel	Rzotkiewicz <i>et al.</i> 2015
<i>Bartonella</i> near <i>henselae</i>	<i>Or. howardi</i>	<i>Glaucomys volans</i>	USA	Reeves <i>et al.</i> 2007a
	<i>Sy. cleopatrae</i>	<i>Gerbillus andersoni allenbyi</i>	Israel	Morick <i>et al.</i> 2010
<i>Bartonella</i> near <i>phoceensis</i>	<i>X. cheopis</i>	<i>R. norvegicus</i> , <i>R. rattus</i>	Egypt	Loftis <i>et al.</i> 2006
<i>Bartonella</i> near <i>quintana</i>	<i>Or. howardi</i>	<i>S. carolinensis</i>	USA	Durden <i>et al.</i> 2004
<i>Bartonella</i> near <i>rochalimae</i>	<i>L. taschenbergi amitina</i>	<i>A. sylvaticus</i>	Spain	Cevidaneš <i>et al.</i> 2017
	<i>X. cheopis</i>	<i>R. norvegicus</i>	Algeria	Bitam <i>et al.</i> 2012
	<i>X. ramesis</i>	ND	Israel	Rzotkiewicz <i>et al.</i> 2015
<i>Bartonella</i> near <i>taylorii</i>	<i>Ct. lushuiensis</i>	<i>Eothenomys</i> sp.	China	Li <i>et al.</i> 2007
<i>Bartonella</i> near <i>tribocorum</i>	<i>X. cheopis</i>	<i>R. rattus</i>	Benin	Leulmi <i>et al.</i> 2014
<i>Bartonella</i> near <i>vinsonii arupensis</i>	<i>Sy. cleopatrae</i>	ND	Israel	Rzotkiewicz <i>et al.</i> 2015
<i>Bartonella</i> sp.	<i>Echinophaga gallinacea</i>	<i>Dipodomys spectabilis</i>	México	Fernández-González <i>et al.</i> 2016
	<i>Ct. andorrensis catalanensis</i>	<i>C. russula</i>	Spain	Cevidaneš <i>et al.</i> 2017
	<i>M. arachis</i>	<i>D. spectabilis</i>	México	Fernández-González <i>et al.</i> 2016
	<i>M. altecpin</i>	<i>D. spectabilis</i> , <i>O. arenicola</i>	México	Fernández-González <i>et al.</i> 2016
	<i>Or. hirsuta</i>	<i>Cynomys</i> sp.	USA	Reeves <i>et al.</i> 2007b
	<i>Sy. cleopatrae</i>	ND	Israel	Rzotkiewicz <i>et al.</i> 2015
	<i>Thrassis aridis</i>	<i>D. spectabilis</i>	México	Fernández-González <i>et al.</i> 2016
	<i>X. cheopis</i>	<i>R. norvegicus</i>	Algeria	Bitam <i>et al.</i> 2012
		<i>R. rattus</i>	Algeria, Israel	Morick <i>et al.</i> 2010; Bitam <i>et al.</i> 2012
<i>R. conorii</i>	<i>Stivalius aporus</i>	<i>Mus caroli</i>	Taiwan	Kuo <i>et al.</i> 2016
<i>R. felis</i>	<i>Acropsylla episema</i>	<i>Apodemus agrarius</i>	Taiwan	Kuo <i>et al.</i> 2016
	<i>Anomiopsyllus nudata</i>	<i>N. albigula</i>	USA	Stevenson <i>et al.</i> 2005
	<i>Ctenocephalides felis</i>	<i>Peromyscus yucatanicus</i>	México	Peniche Lara <i>et al.</i> 2015
		<i>R. norvegicus</i>	Cyprus	Psaroulaki <i>et al.</i> 2006
		<i>R. rattus</i>	Cyprus	Psaroulaki <i>et al.</i> 2006
	<i>Ct. agyrtes</i>	<i>Apodemus flavicollis</i>	Lithuania	Radzijeuskaja <i>et al.</i> 2018
	<i>Ctenophthalmus calceatus calceatus</i>	<i>Lophuromys aquilus</i>	Tanzania	Leulmi <i>et al.</i> 2014
	<i>Ctenophthalmus</i> sp.	<i>R. norvegicus</i>	Portugal	De Sousa <i>et al.</i> 2006
	<i>H. talpae</i>	<i>Micromys minutus</i>	Lithuania	Radzijeuskaja <i>et al.</i> 2018
	<i>L. segnis</i>	<i>Mus</i> sp.	Algeria	Bitam <i>et al.</i> 2009
	<i>Polygenis odiosus</i>	<i>Ototylomys phyllotis</i>	México	Peniche Lara <i>et al.</i> 2015
	<i>S. aporus</i>	<i>M. caroli</i>	Taiwan	Kuo <i>et al.</i> 2016
	<i>X. cheopis</i>	<i>R. norvegicus</i>	Cyprus	Christou <i>et al.</i> 2010
		<i>R. rattus</i>	Cyprus, Madagascar	Christou <i>et al.</i> 2010; Rakotonanahary <i>et al.</i> 2017
		<i>Rattus</i> sp.	Afghanistan, Algeria	Marie <i>et al.</i> 2006; Bitam <i>et al.</i> 2009
<i>R. helvetica</i>	<i>Ct. agyrtes</i>	<i>A. flavicollis</i>	Lithuania	Radzijeuskaja <i>et al.</i> 2018
	<i>M. turbidus</i>	<i>A. flavicollis</i>		
		<i>M. minutus</i>		
	<i>M. walkeri</i>	<i>A. flavicollis</i>		
<i>R. japonica</i>	<i>S. aporus</i>	<i>M. caroli</i>	Taiwan	Kuo <i>et al.</i> 2016
<i>R. monacensis</i>	<i>Ct. agyrtes</i>	<i>A. flavicollis</i>	Lithuania	Radzijeuskaja <i>et al.</i> 2018
<i>R. raoultii</i>	ND	<i>A. flavicollis</i> , <i>Myodes glareolus</i>	Germany	Obiegala <i>et al.</i> 2016
<i>R. typhi</i>	<i>Ctenophthalmus congeneroides</i>	<i>A. agrarius</i>	South Korea	Kim <i>et al.</i> 2010
	<i>L. segnis</i>	<i>R. norvegicus</i>	Cyprus	Christou <i>et al.</i> 2010
		<i>R. rattus</i>	Cyprus, Egypt, Portugal	De Sousa <i>et al.</i> 2006, Loftis <i>et al.</i> 2006; Christou <i>et al.</i> 2010
	<i>Rhadinopsylla insolita</i>	<i>A. agrarius</i>	South Korea	Kim <i>et al.</i> 2010
	<i>Xenopsylla brasiliensis</i>	<i>Mastomys natalensis</i>	Tanzania	Leulmi <i>et al.</i> 2014

Continue...

Bacteria species	Sucking lice	Host	Country	References
		<i>R. rattus</i>	Tanzania	Leulmi et al. 2014
		<i>Rattus</i> sp.	Democratic Republic of the Congo	Leulmi et al. 2014
	<i>X. cheopis</i>	<i>R. norvegicus</i>	Cyprus, Egypt	Loftis et al. 2006; Christou et al. 2010
		<i>R. rattus</i>	Benin, Cyprus, Egypt, Madagascar	Loftis et al. 2006; Christou et al. 2010; Leulmi et al. 2014, Rakotonanahary et al. 2017
		<i>Rattus</i> sp.	Algeria	Bitam et al. 2009
<i>Rickettsia prowazekii</i>	<i>Or. howardii</i>	<i>G. volans</i>	USA	Sonenshine et al. 1978
<i>Candidatus Rickettsia Asemboensis</i>	<i>E. gallinacea</i>	<i>R. rattus</i>	Egypt	Loftis et al. 2006
	<i>S. cleopatrae</i>	ND	Israel	Rzotkiewicz et al. 2015
	<i>X. ramesis</i>	<i>Gerbillus dasyurus</i> , <i>Meriones tristrami</i> , <i>M. musculus</i>	Israel	Rzotkiewicz et al. 2015
<i>Rickettsia felis</i> -like	<i>X. ramesis</i>	ND	Israel	Rzotkiewicz et al. 2015
<i>Rickettsia</i> near <i>monacensis</i>	<i>Oropsylla hirsuta</i>	<i>Cynomys</i> sp.	USA	Reeves et al. 2007b
<i>Rickettsia</i> sp. Oh16	<i>Or. howardi</i>	<i>S. carolinensis</i>	USA	Reeves et al. 2005
<i>Rickettsia</i> sp. TwKM01	<i>S. aporus</i>	<i>A. agrarius</i>	Taiwan	Kuo et al. 2016
<i>Rickettsia</i> endosymbiont of <i>Eucoryphus brunneri</i>	<i>Ct. agyrtes</i>	<i>A. flavicollis</i>	Lithuania	Radzijevska et al. 2018
<i>B. henselae</i>	<i>Neohaematopinus sciuri</i>	<i>S. carolinensis</i>	USA	Durden et al. 2004
<i>B. phoceensis</i>	<i>Hoplopleura pacifica</i>	<i>R. norvegicus</i>	Egypt	Reeves et al. 2006
	<i>Polyplax spinulosa</i>	<i>R. norvegicus</i>	Taiwan	Tsai et al. 2010
	<i>Polyplax</i> sp.	<i>R. rattus</i>	Madagascar	Brook et al. 2017
		<i>Rattus</i> sp.	Thailand	Klangthong et al. 2015
<i>B. rattimassiliensis</i>	<i>Hoplopleura pacifica</i>	<i>R. norvegicus</i>	Egypt	Reeves et al. 2006
	<i>Polyplax spinulosa</i>	<i>R. norvegicus</i>	Egypt, Taiwan	Reeves et al. 2006; Tsai et al. 2010
	<i>Polyplax</i> sp.	<i>R. rattus</i>	Madagascar	Brook et al. 2017
		<i>Rattus</i> sp.	Thailand	Klangthong et al. 2015
<i>B. tribocorum</i>	<i>Polyplax spinulosa</i>	<i>R. norvegicus</i>	Taiwan	Tsai et al. 2010
<i>B. vinsonii</i>	<i>Hoplopleura hirsuta</i>	<i>S. hispidus</i>	México	Sánchez-Montes et al. 2016b
<i>B. washoensis</i>	<i>Neohaematopinus sciuri</i>	<i>S. carolinensis</i>	USA	Durden et al. 2004
<i>Bartonella</i> near <i>tribocorum</i>	<i>Polyplax spinulosa</i>	<i>R. norvegicus</i>	Egypt	Reeves et al. 2006
<i>Bartonella</i> near <i>washoensis</i>	<i>Hoplopleura sciuricola</i>	<i>S. carolinensis</i>	USA	Durden et al. 2004
<i>Bartonella</i> sp.	<i>Polyplax</i> sp.	<i>Thrichomys apereoides</i>	Brazil	Fontalvo et al. 2017
<i>R. prowazekii</i>	<i>Neohaematopinus sciuropteri</i>	<i>G. volans</i>	USA	Sonenshine et al. 1978
	<i>Polyplax spinulosa</i> *	<i>R. norvegicus</i>	México	Mooser et al. 1931
<i>R. typhi</i>	<i>Enderleinellus marmotae</i>	<i>Marmota monax</i>	USA	Reeves et al. 2005
	<i>Hoplopleura pacifica</i>	<i>R. norvegicus</i>	Egypt	Reeves et al. 2006

In México, nine taxa of fleas (*Ctenocephalides felis*, *Maleareus sinomus*, *Meringis parkeri*, *Orchopeas hirsuta*, *O. leucopus*, *O. sexdentatus*, *Pleochaetis exilis*, *Pulex* sp., and *Polygenis odiosus*) and two species of sucking lice (*Hoplopleura hirsuta* and *Polyplax spinulosa*) tested positive for at least one of four validated species of *Bartonella* (*B. vinsonii* and *B. washoensis*) and *Rickettsia* (*R. felis* and *R. prowazekii*). Additionally new lineages of *Bartonella* have been registered in six more flea species (*Echinophaga gallinacea*, *Meringis altipecten*, *M. arachis*, *M. parkeri*, *Pleochaetis exilis*, *Thrassis aridis*, Table 1). These records came from isolated studies carried out in wildlife from the southeast and northern parts, lacking data regarding central México where there is a report of human cases of murine typhus (Centro Nacional de Vigilancia Epidemiológica y Control de Enfermedades

2018; Sánchez-Montes et al. 2019). Additionally, for México, 172 species of fleas and 44 species of sucking lice, have been recorded, then, the inventory of species of both bacteria genera is still far from complete (Sánchez-Montes et al. 2013; Acosta-Gutiérrez 2014).

Due to the great diversity of potential vectors and the historical presence of human cases of murine typhus in the centre of the country; the purpose of this study was to identify the presence and diversity of *Bartonella* and *Rickettsia* species in a focus of murine typhus in Hidalgo, México.

Material and Methods

During August to September 2014, we sampled in two private ranches from Mineral del Monte and Tulancingo de Bravo (Figure 1), in the state of Hidalgo, México, close




Figure 1. Sampling sites along the state of Hidalgo, México. Green: State of Hidalgo; Brown: Huasca de Ocampo; Yellow: Mineral del Monte.

to sites where human murine typhus cases have been reported (CENAPRECE 2016). This study was approved by the Ethics and Research Committee of the Medical Faculty of the Universidad Nacional Autónoma de México [FMED/CI/JMO/004/2012].

In order to identify the presence of several flea-borne and louse-borne pathogens (*Rickettsia* and *Bartonella*) in small mammals and their associated ectoparasites, we trapped small mammals using Sherman traps following Romero-Almaraz *et al.* (2007), under permission FAUT-0170 from the Secretaría del Medio Ambiente y Recursos Naturales. All mammals were sacrificed in accordance with the Guidelines of the American Society of Mammalogists for the Use of Wild Mammals in Research (Sikes *et al.* 2016). We performed the necropsy of each animal, extracting a portion of liver and ear which were fixed in 96 % ethanol until its processing in the laboratory. Additionally, fleas and lice were recovered from host's bodies by manual inspection and fixed in absolute ethanol. Hosts and fleas were identified and deposited at the Mammal Collection and the Flea Collection of the Museo de Zoología "Alfonso L. Herrera" Facultad de Ciencias (MZFC) and Colección del Centro de Medicina Tropical, Facultad de Medicina (CMTFM), both belonging to Universidad Nacional Autónoma de México.

For morphological determination, fleas and lice were mounted on slides using the modified techniques of Kim *et al.* (1986) and Wirth and Marston (1968). Species were identified using specialized taxonomic keys such as Kim *et al.* (1986) for lice and Acosta and Morrone (2003), Has-triter (2004), Hopkins and Rothschild (1971), Morrone *et al.* (2000), and Traub (1950) for fleas.

From collected ectoparasites and hosts tissues, we extracted DNA with the QIAamp® DNA Mini Kit (QIAGEN, Hilden, Germany). As an endogenous internal control and for molecular identification of the ectoparasites, we amplified a

Table 2. Oligonucleotide primers used in this study.

Gen	Primers	Sequence (5'-3')	Length (bp)	Reference
<i>Fleas and lice</i>				
COI (Cytochrome oxidase subunit I)	L6625	CCGGATCCTTYTGRTTYTYGGNCAYCC	400	Hafner <i>et al.</i> 1994
	H7005	CCGGATCCACNACRTARTANGTRTRCRTG		
<i>Rickettsia</i> sp.				
<i>gltA</i> (Citrate synthase)	RpCS.415	GCTATTATGCTTGCGGCTGT	806	de Souza <i>et al.</i> (2006)
	RpCS.1220	TGCATTCTTTCCATTGTGC		
<i>ompB</i> (Outer membrane protein B)	120-M59	CCGCAGGGTTGGTAACTGC	862	Roux and Raoult, 2000
	120-807	CCTTTTAGATTACCGCCTAA		
<i>Bartonella</i> sp.				
<i>gltA</i> (Citrate synthase)	BhCS781.p	GGGGACCAGCTCATGGTGG	379	Norman <i>et al.</i> 1995
	BhCS1137.n	AATGCAAAAAGAACAGTAAACA		

fragment of 400 bp of Cytochrome Oxidase Subunit I (COI) gene. For pathogens detection, we amplified a fragment of *gltA* and *ompB* genes specific for each group using primers and temperature conditions previously reported (Table 2).

The reaction mixture consisted of 12.5 µL of GoTaq® Green Master Mix, 2X of Promega Corporation (Madison, WI, USA), the pair of primers (100 ng each), 6.5 µL nuclease-free water and 30 ng DNA in a final volume of 25 µL (Sánchez-Montes *et al.* 2016a, b).

PCR products were resolved in 2 % agarose gels using TAE buffer at 85 V during 45 minutes and visualized using an ODYSSEY CLx Imaging System (LICOR Biosciences). Purified amplification products were submitted for sequencing at Macrogen Inc., Korea.

Sequences were analysed and edited using Bioedit version 5.0.9 Sequencing Alignment Editor Copyright © program and deposited in GenBank under accession numbers (MG952757 to MG952772). In order to identify the species of *Bartonella* and *Rickettsia*, we used the similarity criteria of the *gltA* and *ompB* genes proposed by La Scola (2003), Fournier and Roullet (2009) and Fournier *et al.* (2003). Global alignments were done using Clustal W (Thompson *et al.* 1994) and the best substitution model was selected based on the lowest BIC (Bayesian Information Criterion) score for each gene using MEGA 6.0 (Tamura *et al.* 2011; Sánchez-Montes *et al.* 2016c). Additionally phylogenetic reconstruction was done using Maximum Likelihood also in MEGA 6.0 and branch support was evaluated over 10,000 bootstrap replications.

Results

We collected 40 rodents from four species (*Mus musculus*, *Peromyscus beatae*, *Rattus norvegicus*, and *Reithrodontomys sumichrasti*), and one shrew (*Sorex ventralis*), which are deposited in the MZFC under the following catalogue numbers LRR001 to LRR040. We detected the presence of *Bartonella* DNA in four samples of liver of two *P. beatae* (2/26 = 7.69 %) and two *R. norvegicus* (2/4 = 50 %). Sequences recovered from *P. beatae* exhibited a similarity of 98 % with *B. vinsonii vinsonii* (a member of the *Bartonella vinsonii* complex) and those from *R. norvegicus* corresponded in a 100 %, respectively with *B. elizabethae* (Figure 2). In the case of




Figure 2. Maximum likelihood (ML) phylogenetic tree generated with *gltA* gene (300 bp) from several members of the genus *Bartonella*. The nucleotide substitution model was the Tamura three parameter model (T92) with discrete Gamma distribution (+G). Bootstrap values higher than 50 are indicated at the nodes. Sequences recovered in the study are marked with blue rhombuses and red triangles.

Rickettsia detection, a single specimen of *R. norvergicus* (1/4 = 25 %) tested positive in samples from liver and ear; we recovered sequences of *gltA* and *ompB* genes which exhibited a similarity of 99 % and 100 % with *R. typhi* (Accession




Figure 3. Maximum likelihood (ML) phylogenetic tree generated with *gltA* and *ompB* genes concatenated (1547 bp) from several members of the genus *Rickettsia*. The nucleotide substitution model was the Tamura three parameter model (T92) with discrete Gamma distribution (+G). Bootstrap values higher than 50 are indicated at the nodes. Sequences recovered in the study are marked with red triangles.

number AE017197) deposited in GenBank (Figure 3). A single *R. norvergicus* specimen presents co-infection between *B. elizabethae* and *R. typhi*.

Hosts were infested by 47 fleas (18 females, 29 males), and 172 sucking lice (60 females, 39 males, 73 nymphs), distributed in six taxa, five species belonging to five families and six genera (Table 3). No fleas or lice were recovered from *M. musculus* and *S. ventralis*. After morphological identification was done, we amplified a fragment of 400 bp of Cytochrome oxidase subunit I (COI) in all ectoparasites recovered, in order to corroborate the identification of all samples, especially of those damaged specimens and nymphal stages. DNA sequences of the COI for four of the six species analysed were deposited in GenBank with the following accession numbers: *C. tecpin* (MG952757), *P. hesperomys adelpha* (MG952758); *P. mathesoni* (MG952759), *P. spinulosa* (MG952772) and *H. reithrodontomydis* (KT151126). No complete sequences were obtained for *J. b. breviloba*. We detected the presence of the same *Bartonella* lineage previously referred in *P. beatae*, in two flea species (six *P. hesperomys adelpha* and 17 *P. mathesoni*) recovered from the two hosts which tested positive and from three others that were negative (Table 3). Sequences from fleas and hosts shape a single cluster within our phylogenetic analysis (Fig. 1). None of the flea or sucking lice species analysed was positive for *Rickettsia* DNA.

Discussion

We report for the first time the presence of two species of *Bartonella* and one of *Rickettsia* in the state of Hidalgo, México. The first *Bartonella* species is a member of the *B. vinsonii* complex, closely related with previous sequences detected in Cricetid rodents and fleas of the northern México (Rubio et al. 2014; Fernández-González et al. 2016). Also, this is the first study to report the presence of a *Bartonella* in the fleas *P. hesperomys adelpha* and *P. mathesoni* and in the host *P. beatae* (Table 1). Our phylogenetic analysis grouped sequences of *B. vinsonii* from *P. hesperomys adelpha*, *P. mathesoni* and *P. beatae* in a single cluster, then, our inference is that both flea species could be the potential vectors of these. Additionally, positive *P. hesperomys adelpha* were recovered from negative hosts, suggesting that these fleas may disseminate the pathogen in non-infected individuals among the rodent population bacteria (Kosoy et al. 1997; Morick et al. 2010). However, it is necessary to carry out tests to verify their vectorial capacity. Both species of fleas have a restricted distribution in México, which extend along the northeastern and central parts of the country, parasitizing several cricetid species such as *Peromyscus levipes*, *P. maniculatus*, *Reithrodontomys megalotis* (*P. mathesoni*) and *P. difficilis* (*P. hesperomys adelpha*), so it is not unexpected that this strain of bacteria is widely distributed in the country (Ponce-Ulloa and Llorente-Bousquets 1993; Hoffman et al. 1989; Whitaker and Morales-Malacara 2005; Acosta and Fernández 2015).

Table 3. Ecological parameters of *Bartonella* and *Rickettsia* species detected in fleas, sucking lice and small mammals in Hidalgo, México.

Host					Ectoparasite											
Family	Species	n	HI	%	BAD	Family	Species	HP	EA	%	A	II	EI	%	BAD	
Ranch 1 Tulancingo de Bravo																
Cricetidae	<i>Peromyscus beatae</i>	20	2	10	<i>Bartonella vinsonii</i>	Ceratophyllidae	<i>Jellisonia breviloba breviloba</i>	2	3	10	0	2	0	0	ND	
							<i>Plusaetis mathesoni</i>	10	27	5	1	3	17	57	<i>Bartonella vinsonii</i>	
							Ctenophtalmidae	<i>Ctenophtalmus tecpin</i>	2	3	10	0	2	0	0	ND
							Leptopsyllidae	<i>Peromyscopsylla hesperomys adelpha</i>	4	7	20	0	2	6	86	<i>Bartonella vinsonii</i>
	<i>Reithrodontomys sumichrasti</i>	2	0	0	ND	Hoplopleuridae	<i>Hoplopleura reithrodontomydis</i>	1	4	50	2	4	0	0	ND	
Soricidae	<i>Sorex ventralis</i>	1	0	0	ND	NR	NR	0	NR	(-)	(-)	(-)	NR	NR	ND	
Ranch 2 Mineral del Monte																
Cricetidae	<i>Peromyscus beatae</i>	6	0	0	ND	Ceratophyllidae	<i>Plusaetis mathesoni</i>	1	3	17	1	3	0	0	ND	
Muridae	<i>Mus musculus</i>	8	0	0	ND	NR	NR	0	NR	(-)	(-)	(-)	NR	NR	ND	
	<i>Rattus norvegicus</i>	4	2	50	<i>Bartonella elizabethae</i>	Polyplacidae	<i>Polyplax spinulosa</i>	4	172	100	43	43	0	0	ND	
		1	25	<i>Rickettsia typhi</i>												

n: Host collected; HI: Number of hosts infected; %: Prevalence; BAD: Bacterial agents detected; HP: Host parasitized; EA: Ectoparasites collected; A: Mean abundance; II: Intensity of infestation; EI: Ectoparasites infected; NR: Not recovered; ND: Not detected.

We also report for the first time the presence of *B. elizabethae* in México, a zoonotic bacterial that may causes endocarditis and neuroretinitis in humans. This agent was reported for the USA in the 1990's, however, is has become an emerging problem in several countries of Southeast Asia, Portugal and France (Regier et al. 2016; Tay et al. 2016). *Bartonella elizabethae* is mainly transmitted by the rat flea *Xenosylla cheopis* (Table 1); however, in our study we did not recovered any fleas from the four *R. norvegicus* analysed. The higher prevalence of *B. elizabethae* in collected murid rodents suggests the presence of this flea or other competent vector in the area (Bitam et al. 2012). Additionally, we compiled evidence for the first time of the presence of *R. typhi* in rodents of the state of Hidalgo. This *Rickettsia* produces febrile cases with a wide range of severity that can lead to systemic failure in less than 5% percent of cases (Zavala-Castro et al. 2009). In the state of Hidalgo, three cases of murine typhus had been reported between 2005 to 2010, nevertheless, in 2015 there was an outbreak with 12 cases (Centro Nacional de Vigilancia Epidemiológica y Control de Enfermedades 2018).

Only one rat reported coinfection by *B. elizabethae* and *R. typhi*, a phenomenon that has been previously reported, probably because both pathogens are transmitted by the same flea species (Table 1). This reinforces the hypothesis of the presence of this vector in the study area (Marie et al. 2006; Bitam et al. 2012; Frye et al. 2015). The presence of positive Norway rats for these two zoonotic pathogens is a risk to human health, because this rodent species invade suburban and urban areas, live and thrive in human settlements and could carry fleas that can feed on human hosts and produce urban outbreaks. Our findings represent the first record of several confirmed zoonotic pathogens that can cause murine typhus and endocarditis in México, which highlight the importance of the establishment of active entomological surveillance in wildlife.

Acknowledgements

We thank to A. Villalpando, O. Escorza and G. Cruz for their help in the logistics and direction of sampling. Additionally to Y. N. Lozano Sardaneta for editing our images. We are indebted to J. C. Sánchez-Montes of the Department for Teaching and Research Branch of the General Directory for Preventive Medicine in Secretaria de Comunicaciones y Transportes, who kindly reviewed our manuscript and provided a number of valuable comments. This work was supported by grants CONACyT 221405 and PAPIIT IN211418. There are no financial or commercial conflicts of interest. Daniel Sokani Sánchez Montes was supported by a fellowship from CONACyT and was a Ph.D. student of Programa de Doctorado en Ciencias Biomédicas, Universidad Nacional Autónoma de México, UNAM.

Literature cited

- ABBOT, P., A. E. AVILES, L. ELLER, AND L. A. DURDEN. 2007. Mixed infections, cryptic diversity, and vector-borne pathogens: evidence from *Polygenis* fleas and *Bartonella* species. Applied and Environmental Microbiology 73:6045–6052.
- ACOSTA, R., AND J. J. MORRONE. 2003. Clave ilustrada para la identificación de los taxones supraespecíficos de Siphonaptera de México. Acta Zoológica Mexicana 89:39–53.
- ACOSTA, R. 2014. Biodiversidad de Siphonaptera en México. Revista Mexicana de Biodiversidad 85:345–352.
- ACOSTA, R., AND J. A. FERNÁNDEZ. 2015. Flea diversity and prevalence on arid-adapted rodents in the Oriental Basin, México. Revista Mexicana de Biodiversidad 86:981–988.
- BITAM, I., B. BAZIZ, T. KERNIF, Z. HARRAT, P. PAROLA, AND D. RAOULT. 2009. Molecular detection of *Rickettsia typhi* and *Rickettsia felis* in fleas from Algeria. Clinical Microbiology and Infection 15:255–256.
- BITAM, I., K. DITTMAR, P. PAROLA, M. F. WHITING, AND D. RAOULT. 2010. Fleas and flea-borne diseases. International Journal of Infectious Diseases 8:667–676.

- BITAM, I., J. M. ROLAIN, V. NICOLAS, Y. L. TSAI, P. PAROLA, V. A. K. B. GUNDI, AND D. RAOULT. 2012. A multi-gene analysis of diversity of *Bartonella* detected in fleas from Algeria. *Comparative Immunology, Microbiology and Infectious Diseases* 35:71–76.
- BREITSCHWERDT, E. B., AND D. L. KORDICK. 2000. Bartonella infection in animals: carriership, reservoir potential, pathogenicity, and zoonotic potential for human infection. *Clinical Microbiology Reviews* 13:428–438.
- BOWN, K. J., M. BENNET, AND M. BEGON. 2004. Flea-borne *Bartonella grahamii* and *Bartonella taylorii* in bank voles. *Emerging Infectious Diseases* 10:684–687.
- BROOK, C. E., Y. BAI, O. YU, C. R. HAFALIANA, S. HAEWON, P. D. ANDREW, C. METCALF, Y. K. MICHAEL, AND D. KATHARINA. 2017. Elucidating transmission dynamics and host-parasite-vector relationships for rodent-borne *Bartonella* spp. in Madagascar. *Epidemics* 25: 56–66.
- CENTRO NACIONAL DE VIGILANCIA EPIDEMIOLÓGICA Y CONTROL DE ENFERMEDADES (CENAPRECE). 2018. Anuarios de Morbilidad durante el periodo 1995–2015, <http://www.epidemiologia.salud.gob.mx/anuario/html/anuarios.html> (accessed 02.03.16.).
- CEVIDANES, A., L. ALTET, A. D. CHIRIFE, T. PROBOSTE, AND J. MILLÁN. 2017. Drivers of *Bartonella* infection in micromammals and their fleas in a Mediterranean peri-urban area. *Veterinary Microbiology* 203:181–188.
- CHRISTOU, C., A. PSAROULAKI, M. ANTONIOU, P. TOUMAZOS, I. IOANNOU, A. MAZERIS, AND Y. TSELENTIS. 2010. *Rickettsia typhi* and *Rickettsia felis* in *Xenopsylla cheopis* and *Leptopsylla segnis* Parasitizing Rats in Cyprus. *The American Journal of Tropical Medicine and Hygiene* 83:1301–1304.
- DE SOUSA, R., P. FOURNIER, M. SILVA, F. AMARO, F. BACELLAR, AND D. RAOULT. 2006. Molecular detection of *Rickettsia felis*, *Rickettsia typhi* and two genotypes closely related to *Bartonella elizabethae*. *The American Journal of Tropical Medicine and Hygiene* 75:727–731.
- DE SOUSA, K. C. M., R. B. AMARAL, H. M. HERRERA, F. M. SANTOS, G. C. MACEDO, P. C. E. ANDRADE, D. M. BARROS, R. Z. MACHADO, AND M. R. ANDRÉ. 2018. Genetic Diversity of *Bartonella* spp. in Wild Mammals and Ectoparasites in Brazilian Pantanal. *Microbiology Ecological* 76:544–554.
- DURDEN, L. A., B. A. ELLIS, C. W. BANKS, J. D. CROWE, AND J. H. OLIVER. 2004. Ectoparasites of gray squirrels in two different habitats and screening of selected ectoparasites for Bartonellae. *Journal of Parasitology* 90:485–489.
- EISEN, R. J., AND K. L. GAGE. 2012. Transmission of flea-borne zoonotic agents. *Annual Review of Entomology* 57:61–82.
- FERNÁNDEZ, A. M., M. Y. KOSOY, A. V. RUBIO, C. B. GRAHAM, J. A. MONTENIERI, L. M. OSIKOWICZ, Y. BAI, R. ACOSTA, R. ÁVILA, K. L. GAGE, AND G. SUZÁN. 2016. Molecular Survey of *Bartonella* Species and *Yersinia pestis* in Rodent Fleas (Siphonaptera) From Chihuahua, México. *Journal of Medical Entomology* 53:199–205.
- FONTALVO, M. C., A. R. FAVACHO, A. C. ARAUJO, N. M. SANTOS, G. M. OLIVEIRA, D. M. AGUIAR, E. R. LEMOS, AND M. C. HORTA. 2017. *Bartonella* species pathogenic for humans infect pets, free-ranging wild mammals and their ectoparasites in the Caatinga biome, Northeastern Brazil: a serological and molecular study. *The Brazilian Journal of Infectious Diseases* S1413–8670, 30316–6.
- FOURNIER, P. E., AND D. RAOULT. 2009. Current knowledge on phylogeny and taxonomy of *Rickettsia* spp. *Annals of the New York Academy of Sciences* 1166:1–11.
- FOURNIER, P. E., J. S. DUMLER, G. GREUB, J. ZHANG, Y. WU, AND D. RAOULT. 2003. Gene sequence-based criteria for identification of new rickettsia isolates and description of *Rickettsia heilongjiangensis* sp nov. *Journal of Clinical Microbiology* 41:5456–5465.
- FRYE, M. J., C. FIRTH, M. BHAT, M. A. FIRTH, X. CHE, D. LEE, AND W. I. LIPKIN. 2015. Preliminary Survey of Ectoparasites and Associated Pathogens from Norway Rats in New York City. *Journal of Medical Entomology* 52:253–259.
- GANNON, W. L., AND ANIMAL CARE AND USE COMMITTEE OF THE AMERICAN SOCIETY OF MAMMALOGISTS. 2016. Guidelines of the American Society of Mammalogists for the use of wild mammals in research and education. *Journal of Mammalogy* 97:663–688.
- GILLESPIE, J. J., N. C. AMMERMAN, M. BEIER, B. S. SOBRAL, AND A. F. AZAD. 2009. Louse- and flea-borne rickettsioses: biological and genomic analyses. *Veterinary Research* 40:1–12.
- HASTRITER, M. W. 2004. Revision of the flea genus *Jellisonia* Traub, 1944 (Siphonaptera: Ceratophyllidae). *Annals of the Carnegie Museum* 73:213–238.
- HOFFMANN, A., M. OJEDA, AND G. LÓPEZ. 1989. Los ectosimbiontes de *Peromyscus difficilis* (J. A. Allen, 1891) (Rodentia:Cricetidae). *Revista de la Sociedad Mexicana de Historia Natural* 40:49–58.
- HOPKINS, G. H. E., AND M. ROTHSCHILD. 1971. An illustred catalogue of the Rothschild Collection of fleas (Siphonaptera) in the British Museum (Nat. Hist.) 5.
- KAEWMONGKOL, G., S. KAEWMONGKOL, H. BURMEJ, M. D. BENNETT, P. A. FLEMING, P. J. ADAMS, A. F. WAYNE, U. RYAN, P. J. IRWIN, AND S. G. FENWICK. 2011. Diversity of *Bartonella* species detected in arthropod vectors from animals in Australia. *Comparative Immunology Microbiology Infection Disease* 34:411–7.
- KAMANI, J., D. MORICK, K. Y. MUMCUOGLU, AND S. HARRUS. 2013. Prevalence and diversity of *Bartonella* species in commensal rodents and ectoparasites from Nigeria, West Africa. *PLoS Negl Tropical Disease* 7:e2246.
- KIM, K. C., H. D. PRATT, AND C. J. STOJANOVICH. 1986. The sucking lice of North America: an illustrated manual for identification. Philadelphia/London: The Pennsylvania State University Press, University Park. Philadelphia, U. S. A.
- KIM, H. C., Y. C. YANG, S. T. CHONG, S. J. KO, S. E. LEE, T. A. KLEIN, AND J. S. CHAE. 2010. Detection of *Rickettsia typhi* and seasonal prevalence of fleas collected from small mammals in the Republic of Korea. *Journal Wildlife Diseases* 46:165–172.
- KLANGTHONG, K., S. PROMSTHAPORN, S. LEEPITAKRAT, A. L. SCHUSTER, P. W. MCCARDLE, M. KOSOY AND R. TAKHAMPUNYA. 2015. The Distribution and Diversity of *Bartonella* Species in Rodents and Their Ectoparasites across Thailand. *PLoS One* 10:e0140856.
- KOSOY, M. Y., R. L. REGNERY, T. TZIANABOS, E. L. MARSTON, D. C. JONES, D. GREEN, G. O. MAUPIN, J. G. OLSON, AND J. E. CHILDS. 1997. Distribution, diversity, and host specificity of *Bartonella* in rodents from the Southeastern United States. *The American Journal of Tropical Medicine and Hygiene* 57:578–88.
- KOSOY, M., D. T. HAYMAN, AND K. S. CHAN. 2012. *Bartonella* bacteria in nature: where does population variability end and a species start. *Infection Genetics and Evolution* 12:894–904.
- KUO, C. C., J. L. HUANG, T. E. LIN, AND H. C. WANG. 2016. Detection of *Rickettsia* Spp. And Host and Habitat Associations of Fleas (Siphonaptera) in Eastern Taiwan. *Medical and Veterinary Entomology* 26:341–350.

- LA SCOLA, B., Z. ZEAITER, A. KHAMIS, AND D. RAOULT. 2003. Gene-sequence-based criteria for species definition in bacteriology: The *Bartonella* paradigm. *Trends Microbiology* 11:318–321.
- LEULMI, H., C. SOCOLOVSKI, A. LAUDISOIT, G. HOUEMENO, B. DAVOUST, I. BITAM, AND P. PAROLA. 2014. Detection of *Rickettsia felis*, *Rickettsia typhi*, *Bartonella* species and *Yersinia pestis* in fleas (Siphonaptera) from Africa. *PLoS Neglected Tropical Diseases* 8:e3152.
- LI, D. M., Q. Y. LIU, D. Z. YU, J. Z. ZHANG, Z. D. GONG AND X. P. SONG. 2007. Phylogenetic analysis of *Bartonella* detected in rodent fleas in Yunnan, China. *Journal of Wildlife Diseases* 43:609–617.
- LIPATOVA, I., A. PAULASKAS, I. PURAITE, J. RADZIEVSKAJA, L. BALCIAUSKAS, AND V. GEDMINAS. 2015. *Bartonella* infection in small mammals and their ectoparasites in Lithuania. *Microbes and Infection* 17:884–888.
- LOFTIS, A. D., W. K. REEVES, D. E. SZUMLAS, M. M. ABBASSY, I. M. HELMY, J. R. MORIARITY, AND G. A. DASCH. 2006. Surveillance of Egyptian fleas for agents of public health significance: *Anaplasma*, *Bartonella*, *Coxiella*, *Ehrlichia*, *Rickettsia*, and *Yersinia pestis*. *The American Journal of Tropical Medicine and Hygiene* 75:41–8.
- MARIE, J. L., P. E. FOURNIER, J. M. ROLAIN, B. DAVOUST, AND D. RAOULT. 2006. Molecular detection of *Bartonella quintana*, *B. elizabethae*, *B. koehlerae*, *B. doshiae*, *B. taylorii*, and *Rickettsia felis* in rodent fleas collected in Kabul, Afghanistan. *The American Journal of Tropical Medicine and Hygiene* 74:436–439.
- MERHEJ, V., E. ANGELAKIS, C. SOCOLOVSKI, AND D. RAOULT. 2014. Genotyping, evolution and epidemiological findings of *Rickettsia* species. *Infection, Genetics and Evolution* 25:122–137.
- MOOSER, H., M. R. CASTANEDA, AND H. ZINSSER. 1931. The transmission of the virus of Mexican typhus from rat to rat by *Polyplax spinulosa*. *Journal of Experimental Medicine* 54:567–575.
- MORICK, D., B. R. KRASNOV, I. S. KHOKHLOVA, G. I. SHENBROT, M. Y. KOSOY, AND S. HARRUS. 2010. *Bartonella* genotypes in fleas (Insecta: Siphonaptera) collected from rodents in the Negev Desert, Israel. *Applied and Environmental Microbiology* 76:6864–6869.
- MORRONE, J. J., R. ACOSTA, AND A. L. GUTIÉRREZ. 2000. Cladistics, biogeography, and host relationships of the flea subgenus *Ctenophthalmus* (*Alloctenus*), with the description of a New Mexican species (Siphonaptera: Ctenophthalmidae). *Journal of the New York Entomological Society* 108:1–12.
- OBIEGALA, A., C. OLTERSDORF, C. SILAGHI, D. KIEFER, M. KIEFER, D. WOLL, AND M. PFEFFER. 2016. *Rickettsia* spp. in small mammals and their parasitizing ectoparasites from Saxony, Germany. *Veterinary Parasitology: Regional Studies and Reports* 5:19–24
- OSIKOWICZ, L. M., S. A. BILLETTER, M. F. RIZZO, M. P. ROOD, A. N. FREEMAN, J. E. BURNS, R. HU, P. JUIENG, V. LOPAREV, AND M. KOSOY. 2016. Distribution and diversity of *Bartonella washoensis* strains in ground squirrels from California and their potential link to human cases. *Vector Borne Zoonotic Diseases* 16:683–690.
- PAROLA, P., O. Y. SANOGO, K. LERDTHUSNEE, Z. ZEAITER, G. CHAUVANCY, J. P. GONZALEZ, R. S. MILLER, S. R. TELFORD, C. WONGSRICHANALAI, AND D. RAOULT. 2003. Identification of *Rickettsia* spp. and *Bartonella* spp. in from the Thai-Myanmar border. *Annals of the New York Academy of Sciences* 990:173–81.
- PENICHE, L. G., R. K. DZUL, O. C. PÉREZ, AND C. J. ZAVALA. 2015. *Rickettsia typhi* in rodents and *R. felis* in fleas in Yucatán as a possible causal agent of undefined febrile cases. *Revista del Instituto de Medicina Tropical de São Paulo* 57:129–132.
- PONCE-ULLOA, H. E., AND J. LLORENTE-BOUSQUETS. 1993. Distribución de los Siphonaptera (Arthropoda, Insecta) en la Sierra de Atoyac de Álvarez, Guerrero, México. México: Universidad Nacional Autónoma de México: Publicaciones especiales del Instituto de Biología, No. 11.
- PSAROULAKI, A., M. ANTONIOU, A. PAPA-EUSTATHIOU, P. TOUMAZOS, P. F. LOUKAIDES, AND Y. TSELENTIS. 2006. Short Report: First Detection of *Rickettsia felis* in *Ctenocephalides felis* fleas parasitizing rats in Cyprus. *The American Journal of Tropical Medicine and Hygiene* 74:120–122.
- RADZIEVSKAJA, J., E. KAMINSKIENĖ, I. LIPATOVA, D. MARDOSAITĖ-BUSAITIENĖ, L. BALČIAUSKAS, M. STANKO, AND A. PAULASKAS. 2018. Prevalence and diversity of *Rickettsia* species in ectoparasites collected from small rodents in Lithuania. *Parasites and Vectors* 11:375–385.
- RAKOTONANAHARY, R. J., A. HARRISON, A. N. MAINA, A. L. RICHARDS, M. RAJERISON, AND S. TELFER. 2017. Molecular and serological evidence of flea-associated typhus group and spotted fever group rickettsial infections in Madagascar. *Parasites and Vectors* 4:125–133.
- REEVES, W. K., M. P. NELDER, AND J. A. KORECKI. 2005. *Bartonella* and *Rickettsia* in fleas and lice from mammals in South Carolina, U. S. A. *Journal Vector Ecology* 30:310–315
- REEVES, W. K., D. E. SZUMLAS, J. R. MORIARITY, A. D. LOFTIS, M. M. ABBASSY, I. M. HELMY, AND G. A. DASCH. 2006. Louse-borne bacterial pathogens in lice (Phthiraptera) of rodents and cattle from Egypt. *Journal of Parasitology* 92:313–318.
- REEVES, W. K., T. E. ROGERS, L. A. DURDEN, AND G. A. DASCH. 2007a. Association of *Bartonella* with the fleas (Siphonaptera) of rodents and bats using molecular techniques. *Journal Vector Ecology* 32:118–22.
- REEVES, W. K., T. E. ROGERS, AND G. A. DASCH. 2007b. *Bartonella* and *Rickettsia* from fleas (Siphonaptera: Ceratophyllidae) of prairie dogs (*Cynomys* spp.) from the western United States. *Journal Parasitology* 93:953–955.
- REGIER, Y., F. O. ROURKE, AND V. A. KEMPF. 2016. *Bartonella* spp. - a chance to establish One Health concepts in veterinary and human medicine. *Parasite Vectors* 9:261–273.
- ROMERO-ALMARAZ, M. L., C. SÁNCHEZ-HERNÁNDEZ, C. GARCÍA-ESTRADA, AND R. D. OWEN. 2007. Mamíferos pequeños. Manual de técnicas de captura, preparación, preservación y estudio. México, Distrito Federal. Las Prensas de Ciencias.
- RUBIO, A. V., R. AVILA-FLORES, L. M. OSIKOWICZ, Y. BAI, G. SUZAN, AND M. Y. KOSOY. 2014. Prevalence and genetic diversity of *Bartonella* strains in rodents from northwestern México. *Vector Borne Zoonotic Disease* 14:838–845.
- RZOTKIEWICZ, S., R. GUTIÉRREZ, B. R. KRASNOV, D. MORICK, I. S. KHOKHLOVA, Y. NACHUM-BIALA, G. BANETH, AND S. HARRUS. 2015. Novel evidence suggests that a '*Rickettsia felis*-like' organism is an endosymbiont of the desert flea, *Xenopsylla ramesis*. *Molecular Ecology* 24:1364–73.
- SÁNCHEZ-MONTES, S., C. GUZMÁN-CORNEJO, L. LEÓN-PANIAGUA, AND G. RIVAS. 2013. A checklist of sucking lice (Insecta: Phthiraptera: Anoplura) associated with Mexican wild mammals, including geographical records and a host-parasite list. *Zootaxa* 3722:183–203.
- SÁNCHEZ-MONTES, S., C. GUZMÁN-CORNEJO, F. RAMÍREZ-CORONA, AND L. LEÓN-PANIAGUA. 2016a. Sucking lice (Insecta: Psocoda:

- Anoplura) associated with rodents distributed in the neotropical region of México. *Revista Mexicana de Biodiversidad* 87:427–435.
- SÁNCHEZ-MONTES, S., C. GUZMÁN-CORNEJO, L. G. HERRERA-MONTALVO, A. D. RICHMAN, J. J. FLORES-MARTÍNEZ, G. F. GARCÍA-RUIZ, M. BERZUNZA-CRUZ, P. GAYTÁN, R. PÉREZ-MONTFORT, AND I. BECKER. 2016b. Presence of *Bartonella* sp. in the sucking louse *Hoplopleura hirsuta* collected from hispid cotton rats (*Sigmodon hispidus*) in México. *Southwestern Entomologist* 41:1031–1036.
- SÁNCHEZ-MONTES, S., C. GUZMÁN-CORNEJO, Y. MARTÍNEZ-NAJERA, I. BECKER, J. M. VENZAL, AND M. B. LABRUNA. 2016c. *Rickettsia lusitaniae* associated with *Ornithodoros yumatensis* (Acari: Argasidae) from two caves in Yucatan, México. *Ticks Tick Borne Disease* 7:1097–1101.
- SÁNCHEZ-MONTES, S., P. COLUNGA-SALAS, E. A. FERNÁNDEZ-FIGUEROA, M. L. HERNÁNDEZ-MEDEL, C. R. BENÍTEZ, AND I. BECKER. 2019. Murine typhus in México City: report of an imported case. *Revista do Instituto de Medicina-Tropical de Sao Paulo*. 61:e16.
- SONENSHINE, D. E., F. M. BOZEMAN, M. S. WILLIAMS, S. A. MASIELLO, D. P. CHADWICK, N. I. STOCKS, D. M. LAUER, AND B. L. ELISBERG. 1978. Epizootiology of epidemic typhus (*Rickettsia prowazekii*) in flying squirrels. *American Journal of Tropical Medicine and Hygiene* 27:339–49.
- STEVENSON, H. L., B. YING, Y. K. MICHAEL, J. A. MONTENIERI, J. L. LOWELL, C., M. C. CHU, K. L. GAGE. 2003. Detection of Novel *Bartonella* Strains and *Yersinia pestis* in Prairie Dogs and Their Fleas (Siphonaptera: Ceratophyllidae and Pulicidae) Using Multiplex Polymerase Chain Reaction. *Journal of Medical Entomology* 40: 329–337.
- TAMURA, K., D. PETERSON, N. PETERSON, G. STECHER, M. NEI, AND S. KUMAR. 2011. MEGA6: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*, <http://dx.doi.org/10.1093/molbev/msr121>.
- TAY, S. T., K. L. KHO, W. Y. WEE, AND S. W. CHOO. 2016. Whole-genome sequence analysis and exploration of the zoonotic potential of a rat-borne *Bartonella elizabethae*. *Acta Tropical* 155:25–33.
- THOMPSON, J. D., D. HIGGINS, AND T. J. GIBSON. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *NucleicAcids Research* 22:4673–4680.
- TRAUB, R. 1950. Siphonaptera of Central America and México: A morphological study of the adeagus with descriptions of new genera and species. *Fieldiana Zoology* 1:1–127.
- TSAI, Y. L., S. T. CHUANG, C. C. CHANG, P. H. KASS, AND B. B. CHOMEL. 2010. *Bartonella* species in small mammals and their ectoparasites in Taiwan. *American Journal of Tropical Medicine and Hygiene* 83:917–923.
- TSAI, Y. L., C. C. CHANG, S. T. CHUANG, AND B. B. CHOMEL. 2011. *Bartonella* species and their ectoparasites: selective host adaptation or strain selection between the vector and the mammalian host? *Comparative Immunology, Microbiology and Infectious Diseases* 34:299–314.
- WHITAKER, J. O., AND J. B. MORALES-MALACARA. 2005. Ectoparasites and other associates (ectodytes) of mammals of México. Pp. 535–666, in *Contribuciones mastozoológicas en homenaje a Bernardo Villa* (Sánchez-Cordero, V., and R. A. Medellín, eds.). Instituto de Biología e Instituto de Ecología, Universidad Nacional Autónoma de México; Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, Distrito Federal, México.
- WINOTO, I. L., H. GOETHERT, I. N. IBRAHIM, I. YUNIHERLINA, C. STOOPS, I. SUSANTI, W. KANIA, J. D. MAGUIRE, M. J. BANGS, S. R. TELFORD 3RD, AND C. WONGSRICHANALAI. 2005. *Bartonella* species in rodents and shrews in the greater Jakarta area. *The Southeast Asian Journal of Tropical Medicine and Public Health* 36:1523–1529.
- WIRTH, W. W., AND N. MARSTON. 1968. A method for mounting small insects on microscope slides in Canada balsam. *Annals of Entomology Society of America* 61:783–784.
- ZAPATA-VALDÉS, C., R. AVILA-FLORES, K. GAGE, J. HOLMES, J. MONTENIERRI, M. KOSOY AND G. SUZÁN. 2018. Mammalian hosts, vectors, and bacterial pathogens in communities with and without black-tailed prairie dogs (*Cynomys ludovicianus*) in Northwestern México. *Journal of Wildlife Diseases* 54:26–33.
- ZAVALA-CASTRO, J. E., J. E. ZAVALA-VELÁZQUEZ, AND J. E. S. UICAB. 2009. Murine typhus in child, Yucatan, México. *Emerging Infectious Diseases* 15:972–974.

Associated editor: Jesús Fernández

Submitted: November 27, 2018; Reviewed: March 18, 2019;

Accepted: April 24, 2019; Published on line: April 30, 2019.